

THE COSTLY BURDEN OF DRUG-RESISTANT TB IN THE U.S.

Multidrug-resistant (MDR) tuberculosis is a major health threat globally. Nearly half a million MDR TB¹ cases are estimated to occur worldwide annually, including cases that are extensively drug-resistant (XDR).²

While MDR and XDRTB are relatively rare in the U.S., their treatment comes at a terrible price – it is very expensive, takes a long time to treat, disrupts lives, and has potentially life-threatening side effects.





Preventing and Controlling MDR and XDR TB in the U.S. Requires:

BETTER
TREATMENT
OPTIONS

RAPID DIAGNOSIS EXPERT TREATMENT OF EVERY TB CASE IMPROVING
GLOBAL TB
DIAGNOSIS AND
TREATMENT

Footnotes

¹ Multidrug-resistant TB is resistant to at least two of the best and most important anti-TB drugs (isoniazid and rifampin).

² Extensively drug-resistant TB is resistant to isoniazid and rifampin among first-line drugs, resistant to any fluoroquinolone and at least one second-line injectable drug.

Source: Marks S et al. Treatment Practices, Outcomes, and Costs of Multidrug-resistant and Extensively Drug-resistant Tuberculosis in the United States. Emerg Infect Dis. 2014;20(5); additional estimates for TB and XDR TB productivity losses due to premature deaths.



Centers for Disease Control and Prevention National Center for HIV/AIDS, Viral Hepatitis, STD, and TR Prevention